

Harveian Oration



Health in an unequal world

Michael Marmot

William Harvey and the Harveian Trust

William Harvey was born in Folkestone on April 1, 1578. He was educated at the King's School, Canterbury, Gonville, and Caius College, Cambridge, and the University of Padua, graduating as doctor of arts and medicine in 1602. He became a Fellow of the Royal College of Physicians in 1607 and was appointed to the Lumleian lectureship in 1615.

In the cycles of his Lumleian lectures over the next 13 years, Harvey developed and refined his ideas about the circulation of the blood. He published his conclusions in 1628 in *Exercitatio Anatomica de Motu Cordis et Sanguinis in Animalibus*, which marks the beginning of clinical science. In it, Harvey considered the structure of the heart, arteries, and veins with their valves. By carefully devised experiments and supported by the demonstration of the unidirectional flow of the blood in the superficial veins of his own forearm, he established that the blood circulated, and did not ebb and flow as had been believed for more than 1000 years.

Harvey was a great benefactor of the College. In 1656 he gave his patrimonial estate of Burmarsh (in Romney Marsh, Kent) to the College to provide for the annual oration and feast. In an indenture dated June 21, 1656, he directed that:

to the end friendship between the members of the said College may be the better continued and maintained, there shall be once every month at the meeting of the Censors at the said College some small collation provided... and once every year there shall be a general feast kept within the said College for all the Fellows that shall please to come...and on the day when such feast shall be kept some one person...shall make an oration...with an exhortation to the Fellows and Members of the said College to search and study out the secret of Nature by way of experiment; and also for the honour of the profession to continue mutual love and affection amongst themselves without which neither the dignity of the College can be preserved nor yet particular men receive that benefit by their admission into the College which else they might expect.

In poor countries, tragically, people die unnecessarily. In rich countries, too, the higher death rate of those in less fortunate social positions is unnecessary. Can there be a link between these two phenomena: inequalities in health between countries and inequalities within? Surely, we could argue, the depredations of grinding poverty— inadequate food, shelter, clean water, and basic medical care or public health—that ravage the lives of the poor in developing countries are different in kind from the way

Professor Sir Michael Marmot

Michael Marmot is Director of the International Institute for Society and Health, and MRC Research Professor of Epidemiology and Public Health at University College London. He has been at the forefront of research into health inequalities for the past 30 years. He is Principal Investigator of the Whitehall Studies of British civil servants, investigating explanations for the striking inverse social gradient in morbidity and mortality. He leads the English Longitudinal Study of Ageing (ELSA) and is engaged in several international research efforts on the social determinants of health. With colleagues, he has a long-term research programme investigating the causes of health disadvantages in the countries of central and eastern Europe and the former Soviet Union. He chairs the Department of Health Scientific Reference Group on tackling health inequalities and the National Institute for Health and Clinical Excellence (NICE) Research and Development Committee. He also chairs the British Heart Foundation (BHF) Primary Prevention Committee and has been a member of the BHF Council since 2000. He was a member of the Royal Commission on Environmental Pollution for 6 years. He chairs the World Cancer Research Fund Panel on Food, Nutrition and Physical Activity and the Prevention of Cancer: a Global Perspective.

In 2000, Michael Marmot received a knighthood for services to epidemiology and understanding health inequalities. Internationally acclaimed, he is a Vice President of the Academia Europaea, a member of the RAND Health Advisory Board, a Foreign Associate Member of the Institute of Medicine, and Chair of the Commission on Social Determinants of Health set up by WHO in 2005. He won the Balzan Prize for Epidemiology in 2004.

This lecture was presented at the Royal College of Physicians, London, UK, on Oct 18, 2006, and is published simultaneously in *Clin Med* 2006; 6: 559–72.

Lancet 2006; 368: 2081–94

Published Online
December 5, 2006
DOI:10.1016/S0140-
6736(06)69746-8



University College London,
London WC1E 6BT, UK
(Prof Sir M Marmot FRCP)

to meet the fundamental human needs of autonomy, empowerment, and human freedom is a potent cause of ill health. In making this case, I shall bring together two rather disparate streams of work. The first is a report of my own research endeavour. I have sought explanations for the social gradient in health, as observed in the Whitehall studies, pointing to the fundamental importance of the circumstances in which people live and work. I emphasise control and the opportunities for full social engagement.¹ The second draws on the work of development economists Amartya Sen and Nicholas Stern. Sen suggests we should see development as freedom to lead the life people have reason to value.² Stern's concept of empowerment is close to Sen's freedom.³ Without empowerment, argues Stern, economic growth will not bring improvements in health and education as well as relief from poverty.

Both in the case of the social gradient in health within countries and the differences in health among countries, changing social conditions to ensure that people have the freedom to lead lives they have reason to value would lead to marked reductions in health inequalities. In both cases the active involvement of individuals, and communities, in decisions that affect their lives is crucial.

Drawing attention to the central role of human freedoms in health is a statement of philosophical position and a call to social action. It is also based on a synthesis of a great deal of medical and social research. William Harvey, who established these orations, famously encouraged the orator to exhort the fellowship of this College to search out the secret of Nature by way of experiment. For an epidemiologist, of course, Nature herself provides the experiments—grand natural experiments. For this epidemiologist, Nature is in league with Society. Societies organise their affairs in different ways and these differences are the grand experiments that provide the researcher with the opportunity to search out the causes of health and disease. I make the connection between social conditions and biological pathways that plausibly provide the link to cardiovascular and other diseases. I do not stop at biology, however.

William Harvey said that the feast and oration should be not only to encourage scientific experiment but for the honour of the profession to continue mutual love and affection among themselves. Outside observers might feel that Fellows of the Royal College of Physicians need no further encouragement to indulge in mutual love. I will focus on the honour of the profession. Our profession seeks not only to understand but also to improve things. Some doctors feel queasy about the prospect of social action to improve health, which smacks of social engineering. Yet, a physician faced with a suffering patient has an obligation to make things better. If she sees 100 patients the obligation extends to all of them. And if a society is making people sick? We have a duty to do what we can to improve the public health and to reduce health inequalities in social groups where these are avoidable

and hence inequitable or unfair. This duty is a moral obligation, a matter of social justice.

Because of this moral obligation, WHO set up a Commission on Social Determinants of Health, which I chair, that is seeking to reach evidence-based policy recommendations on what can be done to reduce health inequalities and improve the health of the disadvantaged.^{4,5} In bringing together research on the social gradient in health within countries and inequalities among countries, I am seeking to provide a clear intellectual justification for the Commission's work.

In arguing that to control disease we need an understanding of disease mechanisms as well as social and political action, I have distinguished antecedents. If an appeal to authority be needed then Rudolph Virchow (1821–1902), whose biology has made appearances in these orations, will amply suffice.

A lesson in the importance of environment

As physicians we are trained that the patient comes first and last. Searching out individual causes of disease, however, does not negate the importance of environmental causes. Studies of migrants show that as environments change, disease rates change. I learnt this lesson through engagement with Leonard Syme and others in a study of men of Japanese ancestry, living in Japan, Hawaii, and California. As Japanese men migrate across the Pacific, the rate of coronary heart disease rises and the rate of stroke falls.⁶ Part of the reason lies with diet and its effect on plasma cholesterol.⁷ But my study, with Len Syme, of Japanese men in California showed a clear relation between degrees of acculturation and coronary heart disease rates—more Americanisation, more disease—that was independent of plasma cholesterol, blood pressure, or smoking.^{8,9} We had some evidence that the particular aspect of Japanese culture that was protective was the degree to which people remained within the protective confines of their ethnic group, thus benefiting from the social cohesion of Japanese culture.

Coming back to Britain, I studied the health of migrants to England and Wales.¹⁰ In general, migrants showed some persisting effects on disease patterns of the country from which they came and clear changes towards the disease patterns of the host country.¹¹ The general point was that as people change their environment, disease rates change. Crucial to this changed environment seemed to be aspects of social relationships, which suggested that the way both to understand disease causation and to change the rates of disease was to pay attention to the social environment.

One thing led to another. Paying attention to the social environment led to the body of research summarised below and to collaboration with other key individuals in scientific enquiry on the relation of society to health.¹² I am now involved in the process of trying to change these relations via the Commission on Social Determinants of Health.

Inequalities in health within societies

Social gradient in richer countries

My starting point is the Whitehall study of British civil servants.¹³ Figure 1 shows results from the 25-year mortality follow-up of men, originally aged 40–69, by age at death. The graded nature of the link between position in the hierarchy and death—the social gradient in mortality—is the challenge to understanding. The gradient is a broader issue than that of poverty and health. We have no difficulty in contemplating how dirty water, poor sanitary facilities, and inadequate nutrition and shelter could cause the diseases of poverty. But Whitehall is not Kibera (the shanty town in Nairobi that is home to 500 000 people). In Whitehall, we have clean water and bathrooms, an excess in supply of calories to eat, and shelter from the elements. Yet among these civil servants, none of whom were destitute, men second from the top of the occupational hierarchy had a higher rate of death than men at the top. Men third from the top had a higher rate of death than those second from the top. I have been writing about this gradient for 28 years¹⁴ and it has been at the core of my research agenda.¹⁵ Why, among men who are not poor in the usual sense of the word, should the risk of dying be intimately related to where they stand in the social hierarchy?

The Whitehall II study, launched 20 years after the first Whitehall study, extended the observation to women.¹⁶ Further, the gradient in mortality extended to most of the major causes of death.¹⁷ Most of our attempts at explanation have focused on cardiovascular disease because there has been such a large body of research on the biological pathways involved in coronary heart disease. The real puzzle is why there should be a social gradient in so many different causes of death.

The gradient is not confined to civil servants. I live and work in the London borough of Camden. In about 25 min I can cycle from Somers Town, just north of University College London, to Hampstead, a little way further north. The life expectancy gap between men living in these two areas is 11 years.¹⁸ I use the ends of the spectrum—the gap between top and bottom—to illustrate the size of the difference but we should always bear in mind that the phenomenon is a gradient: the population is ranged along the spectrum from life expectancy of 70 years for men in Somers Town and St Pancras to 81 years for men in Hampstead.

In Glasgow the gap is bigger. The difference in life expectancy between the most deprived and least deprived areas was 6.9 years in 1981–85; 20 years later this rate had increased to almost 12 years.¹⁹ In and around Washington DC, USA, the gap is bigger still. A 20-year gap exists between poor blacks in downtown Washington and well-off whites in Montgomery County, MD, a short metro ride away.²⁰

The differences in the USA draw attention to the need to clarify what we think lies behind the social gradient in health. Americans have long looked at British society, and their own, and seen that the British class system does not




Figure 1: Mortality over 25 years according to level in the occupational hierarchy. First Whitehall study of British civil servants. Source: Marmot and Shipley.¹³

travel well. Americans do not therefore record their vital statistics, as we have done for so long in the UK, by some measure of social class or socioeconomic group. By contrast, in the USA, statistics are traditionally recorded by race. There has been much discussion as to whether racial differences represent something more than socioeconomic differences;²¹ whether or not the whole story, socioeconomic differences, and the wider social environment are important in generating racial differences.

Robert Erikson used the Swedish Census, linked to mortality, and showed a remarkable social gradient in mortality.²² Men with a PhD had lower mortality rates than those with a master's degree who, in turn, had lower

	Indigenous (men)	Total (men)	Gap (years)
Australia (1996–2001)	59.4	76.6	17.2
Canada (2000)	68.9	76.3	7.4
New Zealand (2000–02)	69.0	76.3	7.3

Sources: Canadian data: Health Canada.²⁵ New Zealand data: New Zealand Life Tables, 2000–02.²⁶ Australian data: Australia Human Rights and Equal Opportunities Commission.²⁷

Table 1: Life expectancy of indigenous peoples in Australia, Canada, and New Zealand.




Figure 2: Under-5 mortality rates per 1000 children by socioeconomic quintile of household. Source: Gwatkin, et al.³⁰

mortality than those with a bachelor's degree, and so on. The social gradient in mortality stretched from top to bottom of the social hierarchy.

The Whitehall and Swedish studies make clear that to understand inequalities in health we need to go beyond binary thinking: poverty bad, non-poverty good. Health follows the social gradient. The challenge is to understand how position in the social hierarchy is related to health. We need to go beyond material deprivation,^{23,24} but recognising the importance of the gradient should not lead us to ignore those at the bottom. Particularly egregious examples of the effects of social exclusion on health come from comparisons of health of indigenous peoples in Canada, New Zealand, and Australia, with that of the total population (table 1).

In each case, the gap between the indigenous group and the total population is substantial. In New Zealand, this gap has been subdivided further by socioeconomic

position. At each social level, Maoris have higher mortality than Europeans at, notionally, the same level.²⁸ In the USA also, Native Americans and Native Alaskans have lower life expectancy than the total population.²⁹

Social gradient in poorer countries

A social gradient in health is observed in many poorer countries. Regrettably, few countries have data systems that allow national data to be disaggregated by some measure of social position; this is especially true for adult mortality. Demographic and health surveys yield data for infant and child deaths (figure 2).³⁰ In each country, the higher the socioeconomic quintile of the household, the lower the rate of child mortality. The result is a gradient, not simply that the poor have high mortality and everyone else is better off.

Although data for adult mortality by social position are sparse in developing countries, data from Matlab in Bangladesh show clearly that increased education is associated with reduced adult mortality.³¹ Similarly, in Chile there are marked differences in adult survival according to education. At age 20 years, women with 13 or more years of education can expect, on average, to live to 72 years of age compared with about 60 years for women with 1–8 years of education.³²

In the transition economies of central and eastern Europe and the former Soviet Union, the social gradient is clear. In the Russian Federation, to take one example, we used a survey method to reconstruct mortality data for men by education. In a population survey we asked whether husbands and brothers were still alive and, if dead, when they died.³³ Figure 3 shows that those with little education had higher mortality than those with university education, and the gap has been growing every year since the collapse of the Soviet Union.




Figure 3: The widening trend in mortality by education in Russia, 1989–2001
45 p 20=probability of living to 65 years when aged 20 years. Source: Murphy, et al.³³

	Life expectancy	
	At birth	GDP
Japan	82.2	27 967
Sweden	80.2	26 750
Switzerland	80.5	30 552
Spain	79.5	22 391
France	79.5	27 677
UK	78.4	27 147
Greece	78.3	19 954
Costa Rica	78.2	9 606
USA	77.4	37 562
Cuba	77.3	5 400
Sri Lanka	74.0	3 778
Russia	65.3	9 230
India	63.3	2 892
Kenya	47.2	1 037
Swaziland	32.5	4 726

The US\$ is taken as the standard, and purchasing power in each country adjusted to it. Source: Human Development Report 2005.³⁴

Table 2: Life expectancy and GDP in purchasing power parity US\$ in 2003

Inequalities between countries

Life expectancy for some countries is shown in table 2, along with gross domestic product (GDP) adjusted for purchasing power. The range of life expectancy is staggeringly large: from 32.5 years (both sexes) in Swaziland to 82 years in Japan.³⁴ This gap in life expectancy has been growing. Figure 4 has much to encourage us and much to depress.³⁴ Life expectancy in the high-income countries of the Organisation for Economic Co-operation and Development (OECD) increased from 71.6 years to 78.8 in the 30 years from 1970–75. In south Asia, the improvement was even more impressive. In two regions, however, the grounds for concern are considerable. In the 1970s, life expectancy in central and eastern Europe and the Soviet Union was 69, ie, 2.6 years behind the high-income OECD countries. 30 years later, the life expectancy had declined to 68.1 years, now 10.7 years behind the OECD countries. At the other end of the life expectancy scale is sub-Saharan Africa where overall life expectancy increased by 0.3 years at a time when it increased by 7.2 in the richest countries. This small increase masks dramatic

declines in many African countries: from 49 to less than 33 in Swaziland; from 50 to 36 in Lesotho; from around 56 to nearly 37 in Zimbabwe.

Some of this decline is due to the terrible toll of HIV/AIDS. Stephen Lewis cries in anguish that we cannot consider AIDS in Africa without considering the state of women and their special vulnerability to rape and sexual violence, early and forced marriage, and lack of access to education, economic and earning power, and rights to own and inherit land or property.³⁵ My statement at the beginning of this lecture that the avoidable deaths of people in poorer countries has to do with the way we organise our affairs in society has no better illustration than the link of sex inequality to the AIDS epidemic in various parts of the world, especially Africa.

Similarly, I do not think we can understand the lack of improvement, even decline, in the countries of central and eastern Europe and the former Soviet Union without considering the opportunity of people to lead flourishing lives in the sense that I am using the term: autonomy and social participation.¹

Inequalities in health are not inevitable

Before diving into the question of explanations of inequalities in health and, hence, what we could do about them, I should make clear that they are not inevitable. Starting first with inequalities among countries, there is nothing inevitable about the data in figure 4. The rapid health improvements in some countries and the lack of improvement in others suggest that changes in social and environmental conditions, and in public health and basic medical care, could do much to change things for the better.

But what of the social gradient in health within countries—is that not inevitable? Social hierarchies are inevitable. Whether or not we could imagine a society without a hierarchy, we would need to search hard to find it. Hierarchies might be inevitable but the health gradient linked to hierarchies is less so. Simple observation shows that the size of the difference in health between top and bottom (as one measure of the size of the gradient) varies within a society over time and among societies.

Figure 3 showed that the gap in expected survival between those with university education and those with little education grew in Russia during the years since the collapse of the Soviet Union. In England and Wales, the gap in male life expectancy between bottom and top social classes grew from 5.5 years to 9.5 years in the space of 20 years (between 1972–76 and 1992–96), and then narrowed slightly.³⁶

Similarly, we see differences in the size of the socioeconomic gaps among countries. Figure 5 shows that the gap in mortality between men in manual and non-manual occupations varies among nine countries in Europe.³⁷ Notably, the gap in Sweden is one of the smallest. The same study showed that relative differences varied somewhat differently.³⁸ Relative differences will depend on




Figure 4: Trends in life expectancy

CEE=Central and Eastern Europe; CIS=Commonwealth of Independent States; OECD=Organisation for Economic Co-operation and Development. Source: Human Development Report.³⁴




Figure 5: Mortality for non-manual and manual workers in nine European countries

Ranked by absolute level of mortality of manual workers; age groups 45–59. Source: Vagero and Erikson.³⁷

the size of the denominator as well as the numerator. In Sweden the mortality rate in non-manual occupations is low and the relative gap is large. The absolute difference is, however, quite small—consistent with what might be expected if Sweden's social democratic policies are leading to smaller inequalities in society.

We need not, then, accept the present size of the social gradient in health as fixed. If it can change, and we can understand why, action is possible to reduce it.

Poverty: lack of money and more

I have set myself the task of trying to achieve a unifying explanation for health inequalities that takes in both the

starting position is that the causes of coronary heart disease or specific cancers will be the same wherever they might be. Causes that apply in rich countries where there has been much research will probably apply to the same diseases in poorer countries, where there has been less research.

Armed with a richer understanding of poverty and the realisation that, increasingly, we need to explain the occurrence of the same disease, we can now turn to the explanatory framework. I want to start by using development economics as an analogy for reduction in health inequalities and then argue that it is more than an analogy.

Nicholas Stern, former Chief Economist at the World Bank, and colleagues argue strongly that development in poor countries rests on two pillars: economic growth and empowerment.³ The factors necessary for economic growth, such as a favourable investment climate, are indeed important. But the effect of growth on poverty, in all its dimensions including health and income, will be far more powerful if it comes hand-in-hand with empowerment of people and communities. Empowerment is both a means to social development and an end in itself if it equates with leading a life one values.

The analogy with health is that we can think of improvement in health of disadvantaged people as built on two intertwining pillars: material conditions for good health, and control of life circumstances or empowerment. In the first category come food, water, sanitation, provision of medical and public health services; in richer countries these will include, among other things, availability of healthy food, opportunities for exercise, and crime-free neighbourhoods. The second pillar is empowerment. Importantly, empowerment could act at the individual level or at the level of the community. As I shall describe below, one way in which empowerment can operate is that control over life circumstances reduces chronic stress and has favourable biological effects. Empowerment at the community level might also be important as a means of securing resources for health. For example, Simon Szreter has made the case that in 19th century England, collective efficacy of communities helped secure access to clean water supplies.⁵¹

The two-pillar (growth and empowerment) model of development could function as more than an analogy. Development, in the sense of relief of poverty, will be important for improvement of health in poor countries. Empowerment might then have an important effect on health through its effect on relief of poverty, as well as more direct effects. Amartya Sen argues that economic growth leads to an improvement in health provided that it is used for poverty reduction and expenditure on public goods. He also points to a second model of health improvement, which he calls the support-led model. The examples of this model are in those communities—Kerala, Costa Rica, Cuba, Sri Lanka—that achieved good health without rapid economic growth. Social cohesion,

which we might think of as empowerment at the community level, seems to play a key part.²

This explanatory framework can then be applied to the social gradient in non-communicable disease. Non-communicable disease is caused by diet, smoking, lack of physical activity, and excess alcohol, among other determinants. But socioeconomic position matters too, not simply because lack of money somehow translates into risk of non-communicable disease. Above a level where material deprivation is no longer the main issue, absolute income is less important than how much one has relative to others. Relative income is important because, as Sen states, it translates into capabilities.⁵² What is important is not so much what you have but what you can do with what you have. Hence control and social engagement.

In rich countries, autonomy and social inclusion might influence disease through their effect on health behaviours such as nutrition, smoking, or alcohol, or through more direct neuroendocrine pathways, ie, chronic stress. These pathways might also operate in poorer countries, but have been less studied. Similarly, at the community level empowerment could lead to better availability of resources for health, or operate through psychosocial processes linked to social capital.

Another feature of the demographic and health scene, common to developed and developing countries, in addition to the predominance of non-communicable diseases, is the ageing of the world's population.^{53,54} For example, the proportion of the population aged over 65 is set to increase by 43% in Italy and 54% in Japan between 2000 and 2030. Yet, in many countries at intermediate stages of development, the projected rate of growth of the ageing population between 2000 and 2030 is much more rapid; for example, 174% increase in India, 227% increase in Mexico, and 277% increase in Malaysia.⁵⁵

Inequalities in health continue in these older age groups. In the English Longitudinal Study of Ageing (ELSA) the onset of disease, disability, and poor cognition happened about 15 years later in people of high social position than in those of lower social position.⁵⁶

Control and social engagement as contributors to inequalities in health

I use control and social engagement as an organising principle, a way of thinking, about human needs that might relate to health on the one hand and the nature of our social arrangements on the other. They are, though, more than an organising principle since there is direct evidence to support them.

My group studied control, initially, in the workplace. The context was the Whitehall II study of British civil servants in which the lower the position in the occupational hierarchy the higher the risk of coronary heart disease and other ailments. The old idea that managers at the top of the hierarchy are under more stress than people below them has been replaced by two

more explanatory models of work stress. The first, the demand control model, posits that stress at work is not caused by how much demand there is, but how much control there is in relation to demand.⁵⁷ The second suggests that imbalance between efforts and rewards is the determinant of chronic stress.⁵⁸ As evolved beings, we are social animals. Part of living in society is expectation of reciprocal rewards—reward in return for effort expended, which is part of what I mean by full social engagement.

We used both of these models in the Whitehall II study and showed that each was related to coronary heart disease risk independent of the other.^{59,60} A review of the evidence showed that there was a high degree of consistency, at least for the control dimension of the demand control model, from other studies.⁶¹ For example, a recent study from the Netherlands showed that beliefs about personal control over life were related to increased risk of coronary heart disease.⁶² As in the Whitehall II study, lack of control was a contributor to the social gradient in coronary heart disease over and above the contribution of the classic risk factors.⁶³

There are ways of depriving people of control over their lives other than in the workplace. In Whitehall II, we also asked people a simple question about how much control they had at home. Women who had less control at home had higher risk of heart disease than women with more control.⁶⁴ We had similar findings for mental illness.⁶⁵

We have had a programme of work investigating the high rates of morbidity and mortality in the countries of central and eastern Europe and the former Soviet Union, of which figure 3 is but one example.^{66,67} In the Czech Republic, as in Whitehall II, low control at work was associated with risk of myocardial infarction and contributed to the social gradient in occurrence of myocardial infarction.⁶⁸ Psychosocial factors at work were also related to depression in the Czech Republic, Poland, and Russia.⁶⁹ We extended the idea of low control over life circumstances beyond the workplace. In a series of cross-sectional studies, we showed that low perceived control was related to poor health.^{70,71} These studies have the drawback that both the predictor (low control) and the outcome (poor health or mental illness) are based on self-reports. There is, then, the possibility simply of biased reporting or contamination of predictor and outcome. In an ecological study, in seven central and eastern European countries, we showed that mean rates of control of a population sample were related to mortality rates of the country from which the sample was drawn.⁷² Contamination of subjectively reported measures is not an issue in this ecological study.

These studies from central and eastern Europe suggest that whole populations can be, more or less, deprived of control over their lives. These communities suffered during the last two decades of the communist period and, in Russia in particular, they suffered when communism and many social institutions collapsed, real incomes

declined by more than half, and there was a dramatic increase in inequality in society.⁶⁷

The other important human need, after autonomy or control, is to be socially engaged. Imbalance between efforts and rewards is part of not having appropriate social reciprocity. I suggest that self-esteem and the esteem of others is part of social engagement. Adam Smith pointed to the importance of having whatever was necessary for taking your place in public without shame.² If I am to include all of these under social engagement it might sound a little nebulous. Two strands of evidence more directly support the link between social engagement and health: social supports and social capital.

Lisa Berkman, in a range of studies and review of the published work, showed that participation in social networks and having a range of social ties is important for health.^{73–75} Sheldon Cohen has contributed to and reviewed this literature with similarly strong conclusions.⁷⁶ Marriage is one obvious domain in which support might be offered or denied. There has been much debate as to the causal interpretation of the health advantage of those who are married. Does health lead to marriage or marriage to health?⁷⁷ I do not propose to reopen that debate here. It is of interest, however, that in Hungary⁷⁸ and the Czech Republic⁷⁹ the rise in mortality during the last two decades of the communist period was more rapid in unmarried men than in married. The increasing disadvantage of the unmarried state was more marked in men than in women, thus adding fuel to the speculation that marriage is more supportive for men than for women.

Taking social networks and supports to the level of the community leads to ideas of social capital—the idea that some communities are marked out by cohesiveness and trust.⁸⁰ The evidence supporting the links between the social capital of a community and health is suggestive. Kawachi has made the idea operational and shown links between social capital and health.⁸¹ Diez-Roux has shown that characteristics of areas affect people's health over and above the socioeconomic characteristics of the individuals who live there.⁸² We have contributed to this literature by showing that these area characteristics might be part of social capital.^{83,84}

My own speculation is that the remarkably good health of Japan could relate to the social cohesion of that society.¹ Similarly, the good health of relatively poor populations such as those of Kerala and Sri Lanka might be attributed to cohesion and social inclusion—particularly of women.²

Pathways linking autonomy and social engagement to health

One set of pathways are those linking autonomy and engagement to resources that lead to better health. Szreter finds no mystery in the link between social capital and health in Victorian England. It led to community action to provide clean water supplies.^{51,85} If social inclusion means

that more people are involved in education, they will benefit from all the economic, social, and psychological benefits that education can bring. As stated above, an approach to controlling HIV/AIDS in southern Africa that ignores the gender disadvantage of women will have little chance of success.

At a more general level, if empowerment is a key strategy for economic and social development then apart from any direct effect of empowerment on health there will be the indirect benefits that accrue from economic and social development of the whole society.

Conversely, if we see autonomy—leading the life one values—as central then resources are important in creating autonomy and social engagement. For example, having a ready supply of potable water, adequate shelter, and bathroom facilities makes leading a life one values more of a possibility.

A second set of pathways relate to the familiar risk factors for chronic disease. There are two questions that relate to my theme: to what extent do the classic risk factors account for the social gradient in disease occurrence; and might differences in health behaviours be one way that autonomy and social engagement change risk of disease?

In the first Whitehall study, a combination of smoking, plasma cholesterol, blood pressure, being overweight, and lack of physical activity accounted for under a third of the social gradient in coronary heart disease mortality.¹⁷ Some estimates put the contribution of smoking higher than that,⁸⁶ but these are based on indirect measures that use lung cancer as a proxy measure for smoking and are higher than some other estimates using proxy measures.⁸⁷ Whatever the precise contribution of these risk factors to the social gradient, they are important. Smoking is now linked with social disadvantage in the developing world as well.⁸⁸

Similarly for obesity, we know that, particularly for women, there is a clear inverse association with socioeconomic position.⁸⁹ This association is now emerging in developing countries. The relation of obesity to education, as a measure of socioeconomic position, is shown in figure 7, with countries classified by degrees of economic development.⁹⁰ Above a GNP per head of about \$2990 (so-called upper middle-income economies), the higher the education the lower the obesity rate.

We need to ask, then, why there should be social gradients in important risk factors for chronic disease. This is not well understood but autonomy and social engagement might be important. The point has been well made that women who are socially disadvantaged and have little opportunity to control their lives or gain personal fulfilment might have little motivation to refrain from smoking.⁹¹

A third type of pathway linking autonomy and engagement, or human flourishing, to health is through chronic stress pathways.^{92,93} Sapolsky has shown the plausibility of stress mechanisms linking social circumstances and status to health in non-human primates.⁹⁴ Both the hypothalamic pituitary adrenal axis and sympatho-adreno-medullary




Figure 7: Women's obesity by quartiles of education

Prevalence ratios based on prevalence of obesity in lowest quartile of education set at 1 for each group of countries. Source: Monteiro, et al.⁹⁰

axes are important. Sapolsky and others have shown that low-status animals have higher cortisol levels than high-status animals.⁹⁵ The strength of this link between low status and cortisol varies across animal species; the more frequently low status is associated with being on the receiving end of stressful encounters the stronger the link between low status and cortisol.⁹⁶

In the Whitehall II study we were particularly interested in two elements of stress pathways linking low social position to increased risk of cardiovascular disease: plasma fibrinogen as an inflammatory marker and the metabolic syndrome. Both are linked to low social position.^{93,97} We have also shown that psychosocial factors are linked to the metabolic syndrome. Figure 8, from the Whitehall II study, shows that the more frequently people reported that their jobs were high strain (low control, high demands, little support) the greater the likelihood of having the metabolic syndrome. Likely mechanisms linking psychosocial factors to the metabolic syndrome are both the autonomic nervous system and hypothalamic pituitary adrenal axis.^{98,99}

My colleague, Andrew Steptoe, studies these psychobiological pathways in the laboratory. He shows the plausibility of linking stressful stimuli to cardiovascular, endocrine, and immune responses,¹⁰¹ and that these




Figure 8: Metabolic syndrome by exposure to iso-strain

Whitehall II study of British civil servants. Odds ratios based on no exposure to iso-strain set at 1, adjusted for age, employment, grade, and health behaviours. Source: Chandola, et al.¹⁰⁰




Figure 9: Differing benefits from government health service expenditure
Average for 21 countries: number of countries where the lowest quintile receives less, the same, or more benefit compared with the highest income quintile.
Source: Yazbeck, et al.¹⁰⁶

responses differ according to socioeconomic position.¹⁰² One interesting finding was that it was not so much the height of the biological stress response that differed by socioeconomic position, but that low-grade civil servants had slower rates of biological recovery after stress.¹⁰³

Medical care?

I have said little about medical care. There is no question that part of improving health in poorer countries, as in richer, is the provision of comprehensive primary care. In a well-organised society there should be universal access to high quality medical care. The whole principle of the UK National Health Service is universal provision and that seems a principle worth exporting. In poorer countries of the world, as in some richer ones, attempts at universal provision do not guarantee universal access. A study in 21 low-income and middle-income countries showed that government expenditure on healthcare favoured, selectively, those of higher incomes (figure 9).¹⁰⁴ In 15 of the countries, people in the top quintile of income obtained the greatest benefit; in only four did people in the bottom quintile of income benefit most; and in two the benefit from government health expenditure was equally shared between top and bottom income groups.




Figure 10: Differences in doctor-diagnosed illness between England and the USA, (55–64-year-olds)
Source: Banks, et al.¹⁰

Gwatkin and colleagues make the point that the health of the poor has more to do with their social and environmental circumstances than with their lack of healthcare. But, since \$380 billion is spent every year on health services in low-income and middle-income countries, it is worth asking whether there are ways to ensure that the expenditure benefits those who need it most—the people at the bottom of the income scale. These people want policies to be not only pro-poor in intent but also in effect.¹⁰⁵

Among rich countries, there is little relation between expenditure on medical care and health. We run the English Longitudinal Study of Ageing (ELSA), which was set up along similar lines to an existing US study, the Health and Retirement Study (HRS). Given that the USA spends \$5274 per head on medical care, and the UK \$2164 (adjusted for purchasing power),³⁴ we were interested in comparing health and health inequalities in the two populations. We compared white men and women, aged 55–64.¹⁰⁷ For each of seven common conditions there is a similar social gradient in health in the two countries, but for each condition more Americans have it than English people. Figure 10 shows the prevalence rates for three of these conditions. One possibility is that Americans have not more illness but more doctors. Greater recognition or recall of illness is not the most likely explanation, however, since in the USA glycosylated haemoglobin was higher, as were levels of C-reactive protein and fibrinogen; levels of high-density lipoprotein cholesterol were lower. Each of these biological markers points to more illness in the USA.

Smoking rates were similar in the USA and England, and alcohol consumption was higher among the English. The prevalence of obesity was about ten percentage points higher in the USA than in England and obesity is likely to be an important factor, particularly for the higher prevalence of diabetes. Although adjusting for the differences in obesity did not abolish the American disadvantage, these were cross-sectional data and we could not look at the cumulative effects of obesity. The higher rate of reported illness is consistent with the higher age-specific mortality rates of the USA compared with the UK in the age range 0–74 years.¹⁰⁸ Interestingly, the USA has lower mortality rates than the UK in those over 74. We do not have the answer to the conundrum posed by the higher rates of illness in the USA but I would speculate that it has, in part, to do with the circumstances in which people live and work—the social determinants of health.

Creating freedom and empowerment

If empowerment is so important for health, how does it arise? Stern conceives of three classes of influence.³ First are individual endowments: assets and human capital. Second are external constraints that come from the context of family, community (including caste and religion), society, and systems of governance, all of which shape people's lives. Third, individuals have internal constraints

on their actions associated with their preferences and perceptions of their role. These classes of influence might be inter-related.

One telling example of societal determinants of empowerment comes from a study of 11–12-year-old children in India.¹⁰⁹ High-caste and low-caste children were given mazes to solve. Despite the high-caste children having higher levels of parental education, the two groups of children did identically on the tests. The tests were then repeated, on different groups, but this time attention was drawn publicly to the caste of the children. Under these circumstances, the lower-caste children did substantially worse. The researchers put this decrement in performance down to an expectation, borne of experience, that lower-caste children would be treated unfairly—it was part of their powerlessness. Confirmation for this speculation was provided by randomly rewarding children, rather than having the decision apparently made by the investigator. Once the rewards were deemed to be fair, the performance of the lower-caste children again matched that of the higher caste.

Power, then, is key.¹¹⁰ Control, autonomy, and freedoms might sound like psychological properties of the individual. As the Stern framework makes clear, however, this is a partial view. Power relations in society, as they operate through social institutions and the opportunities afforded to those in relatively disadvantaged positions, are the social causes of degrees of empowerment. Interestingly, democracy, which should allow more of us some semblance of control, seems to be good for health even after taking other social conditions into account.¹¹¹

An important caution applies to the idea of freedom. In the sense used by Sen, and here, it does not imply privileging the rights of some individuals at the expense of the well-being of others. Human rights can be taken as implying an obligation on society to do what is necessary to bring about the important freedoms that those rights embody.¹¹²

Research as a guide to action

A mark of our civilisation is that we value scientific understanding for its own sake. The Enlightenment brought with it the idea that critical questioning was a better way to the truth than received wisdom. Hence we prize research and scholarship because they enrich us culturally. It is not against this spirit that we might want to apply our knowledge. As physicians we seek not only to understand, but also to make things better. So, too, in public health but the sphere of action is collective rather than individual.¹¹³

We do not seek to improve public health simply by informing individuals of health risks. This is fairly obvious when the health risk comes from the environment. Faced with impure water in a village, we understand that asking each villager to drink expensive bottled water is probably a less effective strategy than piping clean water to the village. This idea might be less obvious when it comes to

behaviours related to chronic disease, but the same insights apply. Individuals choose to drink, smoke, or eat more calories than they consume in physical activity, but their choices are influenced by the environment.

The history of smoking control shows the importance of social action. It has a long history. WHO's Framework Convention on Tobacco Control adopted by the World Health Assembly in 2003 is a landmark achievement in public health.¹¹⁴ Its basic premise is the necessity of governments to be involved in reduction of smoking by individuals.

I have been concerned with the example of alcohol. Evidence shows that the prevalence of heavy drinking is linked to the overall wetness of a population—the total amount of alcohol consumed.^{115,116} Key drivers of alcohol consumption are price and availability.¹¹⁷ Policies to reduce alcohol-related illness, therefore, should deal with price and availability not head off in the opposite direction.¹¹⁸

Social action is as important when it comes to autonomy and opportunities for full social engagement. I was distressed to learn that, at a counselling session for workers who were about to lose their jobs, they were told that Professor Marmot's research shows that control over your own lives is good for health. Now that they would not have to come into work every day these about-to-be-unemployed individuals could look forward to taking control over their lives.

This, of course, is a caricature of the research findings, even a travesty. Individuals' opportunities to control their lives, to be empowered, and to participate fully in society are heavily determined by the way we organise our affairs in society. An excellent review considered the ability of empowerment to improve health.¹¹⁹ It provides encouragement that social change based on insight and understanding can lead to greater empowerment. The review cautions that evidence for the direct effect of improvements in empowerment on health is more limited.

It was precisely to marshal the evidence on social action to improve health that the WHO set up the Commission on Social Determinants of Health. We have set up nine "knowledge networks": early child development, employment conditions, social exclusion, women and gender equity, urban settlements, globalisation, health systems, priority public health conditions, and a cross-cutting network on measurement and evidence. We have a number of other mechanisms for generating the knowledge needed for action: distilling the experience of countries that have been taking action, working with civil society groups, engaging with other organisations.¹²⁰ The Commission is due to report in 2008. Its driving principle is social justice: to reduce unfair differences in health between social groups within a country and between countries. A key mechanism is evidence-based policy. Evidence itself is not enough. There has to be the desire, the political will, for change. Given that will—a big given but I am an optimist—the evidence of what works will be a great help.

The physician and social change

Rudolph Virchow has featured many times in these Lectures. Paul Nurse, for example, quotes Virchow's understanding of cells: "that every animal appears as a sum of vital units, each of which bears in itself the complete characteristics of life". My first contact with Virchow's writing was in relation to his studies of the blood and blood vessels that are important still for our understanding of the pathology of atherosclerosis.

As well as being a scientist who contributed so much to our understanding of pathology, Virchow was also concerned with improving the public health. He wrote: "If medicine is to fulfil her great task, then she must enter the political and social life. Do we not always find the diseases of the populace traceable to defects in society?" He went on: "If disease is an expression of individual life under unfavourable circumstances, then epidemics must be indicative of mass disturbances".^{121,122}

Since disease so often results from poverty, he said, then physicians are the "natural attorneys of the poor", and social problems should largely be solved by them.

I have made the case that a richer understanding of poverty, based on control and social engagement, links the social gradient in health, and poverty and health. We should focus not only on extremes of income poverty but on the opportunity, empowerment, security, and dignity that disadvantaged people want in rich and poor countries alike.

For Virchow, then, it was not biology or society, but both. We need biological understanding of disease but we need, too, understanding of how society influences biology, in order to change disease risk. This social understanding is central to the process of change to reduce the burden of disease.

As physicians we need to be the natural attorneys of the disadvantaged. The Commission on Social Determinants of Health was launched in Chile. It seemed appropriate, then and now, to quote Chile's Pablo Neruda and invite you to: "rise up with me...against the organisation of misery."¹²³

References

- Marmot M. Status syndrome. London: Bloomsbury, 2004.
- Sen A. Development as freedom. New York: Alfred A Knopf, 1999.
- Stern N, Dethier J-J, Rogers H. Growth and empowerment: making development happen. Cambridge, Mass: MIT Press, 2004.
- Marmot M. Social determinants of health inequalities. *Lancet* 2005; **365**: 1099–104.
- Lee JW. Public health is a social issue. *Lancet* 2005; **365**: 1005–06.
- Gordon T. Further mortality experience among Japanese Americans. *Public Health Report* 1967; **82**: 973–84.
- Keys A, Kimura N, Kuzukawa A, et al. Lessons from serum cholesterol studies in Japan, Hawaii and Los Angeles. *Ann Intern Med* 1958; **48**: 83–94.
- Marmot MG, Syme SL, Kagan A, et al. Epidemiologic studies of coronary heart disease and stroke in Japanese men living in Japan, Hawaii and California: Prevalence of coronary and hypertensive heart disease and associated risk factors. *Am J Epidemiol* 1975; **102**: 514–25.
- Marmot MG, Syme SL. Acculturation and coronary heart disease in Japanese Americans. *Am J Epidemiol* 1976; **104**: 225–47.
- Marmot MG, Adelstein AM, Bulusu L. Immigrant mortality in England and Wales 1970–78. London: HMSO, 1984.
- Marmot MG, Adelstein AM, Bulusu L. Lessons from the study of immigrant mortality. *Lancet* 1984; **1**: 1455–58.
- Marmot M, Wilkinson RG, eds. Social determinants of health, 2nd edn. Oxford: Oxford University Press, 2005.
- Marmot MG, Shipley MJ. Do socioeconomic differences in mortality persist after retirement? 25 year follow up of civil servants from the first Whitehall study. *BMJ* 1996; **313**: 1177–80.
- Marmot MG, Rose G, Shipley M, Hamilton PJS. Employment grade and coronary heart disease in British civil servants. *J Epidemiol Community Health* 1978; **32**: 244–49.
- Marmot MG. Status syndrome: a challenge to medicine. *JAMA* 2006; **295**: 1304–07.
- Marmot MG, Davey Smith G, Stansfeld SA, et al. Health inequalities among British Civil Servants: the Whitehall II study. *Lancet* 1991; **337**: 1387–93.
- van Rossum CTM, Shipley MJ, Van de Mheen H, et al. Employment grade differences in cause specific mortality. A 25 year follow up of civil servants from the first Whitehall study. *J Epidemiol Community Health* 2000; **54**: 178–84.
- London Health Observatory. Life expectancy at birth, 2000–2004, Ward Level. www.lho.org.uk (accessed July 24, 2006).
- Hanlon P, Walsh D, Whyte B. Let Glasgow flourish. Glasgow: Glasgow Centre for Population Health, 2006.
- Murray CJL, Michaud CM, McKenna MT, Marks JS. US patterns of mortality by county and race: 1965–94. Cambridge, MA: Harvard Center for Population and Development Studies, 1998: 1–97.
- Williams DR. Race, socioeconomic status and health. The added effects of racism and discrimination. In: Marmot M, McEwen BS, Adler NE, Stewart J, eds. Socioeconomic status and health in industrial nations. Social and biological pathways. Annals of the New York Academy of Sciences, Vol 896. New York: New York Academy of Sciences, 1999: 173–88.
- Erikson R. Why do graduates live longer? In: Jonsson JO, Mills C (eds), Cradle to grave: Life-course change in modern Sweden. Durham: Sociology Press, 2001.
- Lynch JW, Davey-Smith G, Kaplan GA, House JS. Income inequality and mortality: importance to health of individual income, psychosocial environment, or material conditions. *BMJ* 2000; **320**: 1200–04.
- Marmot M, Wilkinson RG. Psychosocial and material pathways in the relation between income and health: a response to Lynch et al. *BMJ* 2001; **322**: 1233–36.
- Health Canada. A statistical profile on the health of First Nations in Canada for the year 2000. http://www.hc-sc.gc.ca/fnih-spni/pubs/gen/stats_profil_e.html (accessed July 26, 2006).
- New Zealand. New Zealand Life Tables 2000–2002. <http://www.stats.govt.nz/analytical-reports/nz-life-tables-2000-2002/appendix1.htm> (accessed July 26, 2006).
- Human Rights and Equal Opportunity Commission. A statistical overview of Aboriginal and Torres Strait Islander peoples in Australia 2003. http://www.humanrights.gov.au/social_justice/statistics/index.html (accessed July 26, 2006).
- Tobias MI, Cheung J. Monitoring health inequalities: life expectancy and small area deprivation in New Zealand. *Popul Health Metrics* 2003; **1**: 2.
- Bramley D, Hebert P, Tuzzio L, Chassin M. Disparities in indigenous health: a cross-country comparison between New Zealand and the United States. *Am J Public Health* 2005; **95**: 844–50.
- Gwatkin D, Johnson K, Wagstaff A, Rutstein S, Pande R. Socio-economic differences in health, nutrition, and population—45 countries. World Bank; 2000. <http://www.worldbank.org/prem/poverty/health/data/statusind.htm> (accessed Aug 16, 2006).
- Hurt LS, Ronsmans C, Saha S. Effects of education and other socioeconomic factors on middle age mortality in rural Bangladesh. *J Epidemiol Community Health* 2004; **58**: 315–20.
- Vega J, Hollstein RD, Delgado I, et al. Socioeconomic health inequities in an intermediate-development nation: Chile, 1985–1996. In: Evans T, Whitehead M, Diderichsen F, Bhuiya A, Wirth M (eds). Challenging health inequities: from ethics to action. New York: Oxford University Press, 2001.
- Murphy M, Bobak M, Nicholson A, Rose R, Marmot M. The widening gap in mortality by educational level in the Russian Federation, 1980–2001. *Am J Public Health* 2006; **96**: 1293–99.

- 34 United Nations Development Programme. Human Development Report 2005: International cooperation at a crossroads. Aid, trade and security in an unequal world. New York: UNDP, 2005.
- 35 Lewis S. Race against time. Toronto, ON, Canada: House of Anansi Press, 2005.
- 36 Donkin A, Goldblatt P, Lynch K. Inequalities in life expectancy by social class, 1972–1999. *Health Stat Q* 2002; **15**: 5–15.
- 37 Vagero D, Erikson R. Socioeconomic inequalities in morbidity and mortality in Western Europe. *Lancet* 1997; **350**: 516–18.
- 38 Mackenbach JP, Bos V, Andersen O, et al. Widening socioeconomic inequalities in mortality in six Western European countries. *Int J Epidemiol* 2003; **32**: 830–37.
- 39 Preston SH. The changing relation between mortality and level economic development *Bull WHO* 2003; **81** (11): 833–41. (Reprinted from *Population Studies* 1975; **29**: 231–48.)
- 40 World Bank. World Development Report 1993. New York: Oxford University Press, 1993.
- 41 United Nations Development Group. Millennium Development Goals, 2000. www.developmentgoals.org.
- 42 Sachs JD, McArthur JW. The Millennium Project: a plan for meeting the Millennium Development Goals. *Lancet* 2005; **365**: 347–53.
- 43 Human Rights and Equal Opportunity Commission. A statistical overview of Aboriginal and Torres Island Strait peoples in Australia 2006. www.hreoc.gov.au/social_justice/statistics/index.html (accessed July 27, 2006).
- 44 Deaton A. Health in an age of globalization. *Brookings Trade Forum* 2004; **2004**: 83–130.
- 45 Narayan D, Patel R, Schafft K, Rademacher A, Koch-Schulte S. Voices of the poor: can anyone hear us? New York: Oxford University Press for the World Bank, 2000.
- 46 Horton R. Rediscovering human dignity. *Lancet* 2004; **364**: 1081–85.
- 47 Marmot M. Dignity and inequality. *Lancet* 2004; **364**: 1019–21.
- 48 Gordon D, Townsend P (eds). *Breadline Europe*. Bristol: Policy Press, 2000.
- 49 Department of Health. Tackling health inequalities: status report on the programme for action. London: Department of Health Publications, 2005.
- 50 WHO. Preventing chronic diseases: a vital investment. Geneva: World Health Organization, 2005.
- 51 Szreter S. Health and wealth: studies in history and policy. New York: University of Rochester Press, 2005.
- 52 Sen A. *Inequality reexamined*. Oxford: Oxford University Press, 1992.
- 53 Panel on a Research Agenda and New Data for an Aging World, Committee on Population, Committee on National Statistics. Preparing for an aging world. Washington: National Academy Press, 2001.
- 54 Vaupel JW, Carey JR, Christensen K, et al. Biodemographic trajectories of longevity. *Science* 1998; **280**: 855–60.
- 55 Kinsella K, Velkoff VA, US Census Bureau. An aging world: 2001. Washington DC: US Government Printing Office, 2001.
- 56 Marmot M, Banks J, Blundell R, Lessof C, Nazroo J. Health, wealth and lifestyles of the older population in England. The 2002 English Longitudinal Study of Ageing. London: Institute for Fiscal Studies, 2003.
- 57 Karasek R, Theorell T. *Healthy work: stress, productivity, and the reconstruction of working life*. New York: Basic Books, 1990.
- 58 Siegrist J, Marmot M. Social inequalities in health: the basic facts. In: Siegrist J, Marmot M (eds). *Social inequalities in health*. New York: Oxford University Press, 2006: 1–25.
- 59 Bosma H, Peter R, Siegrist J, Marmot MG. Two alternative job stress models and the risk of coronary heart disease. *Am J Publ Health* 1998; **88**: 68–74.
- 60 Kuper H, Singh-Manoux A, Siegrist J, Marmot M. When reciprocity fails: effort-reward imbalance in relation to coronary heart disease and health functioning within the Whitehall II Study. *Occup Environ Med* 2002; **59**: 777–84.
- 61 Kuper H, Marmot M, Hemingway H. Systematic review of prospective cohort studies of psychosocial factors in the etiology and prognosis of coronary heart disease. *Semin Vasc Med* 2002; **2**: 267–314.
- 62 Bosma H, Van Jaarsveld CHM, Tuinstra J, et al. Low control beliefs, classical coronary risk factors, and socio-economic differences in heart disease in older persons. *Soc Sci Med* 2005 Feb; **60**: 737–45.
- 63 Marmot MG, Bosma H, Hemingway H, Brunner E, Stansfeld S. Contribution of job control and other risk factors to social variations in coronary heart disease incidence. *Lancet* 1997; **350**: 235–9.
- 64 Chandola T, Kuper H, Singh-Manoux A, Bartley M, Marmot M. The effect of control at home on CHD events in the Whitehall II study: gender differences in psychosocial domestic pathways to social inequalities in CHD. *Soc Sci Med* 2004; **58**: 1501–9.
- 65 Griffin J, Fuhrer R, Stansfeld SA, Marmot MG. The importance of low control at work and home on depression and anxiety: do these effects vary by gender and social class? *Soc Sci Med* 2002; **54** (5): 783–98.
- 66 Bobak M, Marmot MG. East-West mortality divide and its potential explanations: proposed research agenda. *BMJ* 1996; **312**: 421–5.
- 67 Marmot M, Bobak M. International comparators and poverty and health in Europe. *BMJ* 2000; **321**: 1124–8.
- 68 Bobak M, Hertzman C, Skodova Z, Marmot M. Association between psychosocial factors at work and non-fatal myocardial infarction in a population based case-control study in Czech men. *Epidemiology* 1998; **9**: 43–7.
- 69 Pikhart H, Bobak M, Pajak A, et al. Psychosocial factors at work and depression in three countries of Central and Eastern Europe. *Soc Sci Med* 2004 Apr; **58**: 1475–82.
- 70 Bobak M, Pikhart H, Hertzman C, Rose R, Marmot M. Socioeconomic factors, perceived control and self-reported health in Russia. a cross-sectional survey. *Soc Sci Med* 1998; **47**: 269–79.
- 71 Bobak M, Pikhart H, Rose R, Hertzman C, Marmot M. Socioeconomic factors, material inequalities, and perceived control in self-rated health: cross-sectional data from seven post-communist countries. *Soc Sci Med* 2000; **51**: 1343–50.
- 72 Pikhart H. *Social and psychosocial determinants of self-rated health in Central and Eastern Europe*. Boston: Kluwer Academic Publishers, 2002.
- 73 Berkman LF, Glass T. Social integration, social networks, social support, and health. In: Berkman LF, Kawachi I (eds). *Social epidemiology*. New York: Oxford University Press, 2000: 137–73.
- 74 Berkman LF, Glass T, Brissette I, Seeman TE. From social integration to health: Durkheim in the new millennium. *Soc Sci Med* 2000; **51**: 843–57.
- 75 Berkman LF, Melchior M, Chastang J-F, et al. Social integration and mortality: a prospective study of French employees of electricity of France-Gas of France. *Am J Epidemiol* 2004; **159**: 167–74.
- 76 Cohen S, Gottlieb BH, Underwood LG. Social relationships and health. In: Cohen S, Underwood LG, Gottlieb BH (eds). *Social support measurement and intervention: a guide for health and social scientists*. New York: Oxford University Press, 2000: 3–24.
- 77 Goldman N. Marriage selection and mortality patterns: inferences and fallacies. *Demography* 1993; **30**: 189–208.
- 78 Hajdu P, McKee M, Bojan F. Changes in premature mortality differentials by marital status in Hungary and in England and Wales. *Eur J Pub Health* 1995; **5**: 259–64.
- 79 Blazek J, Dzurova D. The decline of mortality in the Czech Republic during the transition: a counterfactual case study. In: Cornia GA, Paniccia R (eds). *The mortality crisis in transitional economies*. New York: Oxford University Press, 2000: 303–27.
- 80 Putnam RD. *Bowling alone: America's declining social capital*. *J Democracy* 1995; **6**: 65–78.
- 81 Kawachi I, Kennedy BP. *The health of nations: why inequality is harmful to your health*. New York: New Press, 2002.
- 82 Diez Roux AV, Merkin SS, Arnett D, et al. Neighborhood of residence and incidence of coronary heart disease. *N Engl J Med* 2001; **345**: 99–106.
- 83 Stafford M, Bartley M, Boreham R, et al. Neighbourhood social cohesion and health: investigating associations and possible mechanisms. In: Morgan A, Swann C (eds). *Social capital and health. Issues of definition, measurement and links to health*. London: Health Development Agency, 2004: 111–31.
- 84 Stafford M, Bartley M, Wilkinson R, et al. Measuring the social environment: social cohesion and material deprivation in English and Scottish neighbourhoods. *Environ Planning A* 2003; **35**: 1459–75.

- 85 Szreter S, Woolcock M. Health by association? Social capital, social theory, and the political economy of public health. *Int J Epidemiol* 2004; **33**: 650–67.
- 86 Jha P, Peto R, Zatonski W, et al. Social inequalities in male mortality, and in male mortality from smoking: indirect estimation from national death rates in England and Wales, Poland, and North America. *Lancet* 2006; **368**: 367–70.
- 87 Marmot M. Smoking and inequalities. *Lancet* 2006; **368**: 341–42.
- 88 Bobak M, Jha P, Nguyen S, Jarvis M. Poverty and smoking. In: Jha P, Chaloupka K (eds). Tobacco control in developing countries. New York: Oxford University Press, 2000: 42–61.
- 89 English Longitudinal Study of Ageing. Retirement, health and relationships of the older population in England: the 2004 English Longitudinal Study of Ageing, Wave 2. London: Institute for Fiscal Studies, 2006.
- 90 Monteiro CA, Conde WL, Lu B, Popkin BM. Obesity and inequities in health in the developing world. *Int J Obes Relat Metab Disord* 2004; **28**: 1181–86.
- 91 Graham H. Hardship and health in women's lives. London: Harvester Wheatsheaf, 1993.
- 92 Brunner EJ. Toward a new social biology. In: Berkman LF, Kawachi I (eds). Social epidemiology. New York: Oxford University Press, 2000.
- 93 Brunner EJ, Juneja M, Marmot MG. Abdominal obesity and disease are linked to social position. *BMJ* 1998; **316**: 508–09.
- 94 Sapolsky RM. Why zebras don't get ulcers: an updated guide to stress, stress-related diseases, and coping. New York: WH Freeman, 1998.
- 95 Sapolsky R. The physiology and pathophysiology of unhappiness. In: Kahneman D, Diener E, Schwarz N (eds). Well-being: the foundations of hedonic psychology. New York: Russell Sage Foundation, 1999: 453–69.
- 96 Abbott DH, Keverne EB, Bercovitch FB, et al. Are subordinates always stressed? A comparative analysis of rank differences in cortisol levels among primates. *Horm Behav* 2003; **43**: 67–82.
- 97 Brunner EJ, Davey Smith G, Marmot MG, et al. Childhood social circumstances and psychosocial and behavioural factors as determinants of plasma fibrinogen. *Lancet* 1996; **347**: 1008–13.
- 98 Brunner EJ, Hemingway H, Walker BR, et al. Adrenocortical, autonomic, and inflammatory causes of the metabolic syndrome. *Circulation* 2002; **106**: 2659–65.
- 99 Hemingway H, Shipley M, Brunner E, et al. Does autonomic function link social position to coronary risk? The Whitehall II Study. *Circulation* 2005; **111**: 3071–77.
- 100 Chandola T, Brunner E, Marmot M. Chronic stress at work and the metabolic syndrome: prospective study. *BMJ* 2006; **332**: 521–5.
- 101 Steptoe A. Psychobiological processes linking socio-economic position with health. In: Siegrist J, Marmot M (eds). Social inequalities in health: new evidence and policy implications. Oxford: Oxford University Press, 2006: 101–26.
- 102 Steptoe A, Marmot M. The role of psychobiological pathways in socio-economic inequalities in cardiovascular disease risk. *Eur Heart J* 2002; **23**: 13–25.
- 103 Steptoe A, Kunz-Ebrecht S, Rumley A, Lowe GD. Prolonged elevations in haemostatic and rheological responses following psychological stress in low socioeconomic status men and women. *Thromb Haemost* 2003; **89**: 83–90.
- 104 Gwatkin DR, Wagstaff A, Yazbeck AS (eds). Reaching the poor with health, nutrition, and population services. Washington DC: World Bank, 2005.
- 105 Marmot M. Health inequalities: good intentions and good results? *Lancet* 2006; **367**: 201–2.
- 106 Yazbeck AS, Gwatkin DR, Wagstaff A, Qamruddin J. Why were the Reaching the Poor Studies undertaken? In: Gwatkin DR, Wagstaff A, Yazbeck AS (eds). Reaching the poor: with health, nutrition, and population services. Washington DC: World Bank, 2005: 3–26.
- 107 Banks J, Marmot M, Oldfield Z, Smith JP. Disease and disadvantage in the United States and in England. *JAMA* 2006; **295**: 2037–45.
- 108 Jarman B, Aylin P. Death rates in England and Wales and the United States: variation with age, sex, and race. *BMJ* 2004; **329**: 1367.
- 109 Hoff K, Pandey P. Belief systems and durable inequities: an experimental investigation of Indian caste. World Bank Policy Research Working Paper. Report No. 3351. Washington DC: World Bank, 2004.
- 110 Farmer P. Pathologies of power: health, human rights, and the new war on the poor. Berkeley: University of California Press, 2003.
- 111 Besley T, Kudamatsu M. Health and democracy. *Am Econ Rev* 2006; **96**: 313–18.
- 112 Sen A. Elements of a theory of human rights. *Philos Public Aff* 2004; **32**: 315–56.
- 113 Beaglehole R, Bonita R, Horton R, Adams O, McKee M. Public health in the new era: improving health through collective action. *Lancet* 2004; **363**: 2084–86.
- 114 Yach D, Hawkes C, Epping-Jordan JE, Galbraith S. The World Health Organization's Framework Convention on Tobacco Control: implications for global epidemics of food-related deaths and disease. *J Public Health Policy* 2003; **24**: 274–90.
- 115 Edwards G, Anderson P, Babor TF, et al. Alcohol policy and the public good. New York: Oxford University Press, 1994.
- 116 Colhoun H, Ben-Shlomo Y, Dong W, Bost L, Marmot M. Ecological analysis of collectivity of alcohol consumption in England: importance of average drinker. *BMJ* 1997; **314**: 1164–68.
- 117 Academy of Medical Sciences. Calling time: the nation's drinking as a major health issue. London: AMS, 2004.
- 118 Marmot M. Evidence based policy or policy based evidence? Editorial. *BMJ* 2004; **328**: 906–7.
- 119 Wallerstein N. What is the evidence on effectiveness of empowerment to improve health? Copenhagen: WHO Regional Office for Europe (Health Evidence Network Report). www.euro.who.int/Document/E88086.pdf (accessed Feb 1, 2006).
- 120 Commission on Social Determinants of Health. 2005. http://www.who.int/social_determinants (accessed July 26, 2006).
- 121 DeWatt DA, Pincus T. The legacies of Rudolf Virchow: Cellular medicine in the 20th century and social medicine in the 21st century. *IMAJ* 2003; **5**: 395–97.
- 122 Friedlander E. Rudolf Virchow on pathology education. <http://www.pathguy.com/virchow.htm> (accessed July 26, 2006).
- 123 Neruda P. The captain's verses. London: Anvil Press, 1994.